

## Evaluation of myocardial perfusion and function after kidney transplantation by Gated SPECT myocardial perfusion scintigraphy

Armaghan Fard-Esfahani<sup>1</sup>, Babak Fallahi<sup>1</sup>, Sahar Mirpour<sup>1</sup>, Ali Gholamrezanezhad<sup>1</sup>,  
Ezatollah Abdi<sup>2</sup>, Mohammad Karimi<sup>1</sup>, Davood Beiki<sup>1</sup>, Alireza Emami-Ardekani<sup>1</sup>,  
Fariba Akhzari<sup>3</sup>, Mojtaba Ansari<sup>4</sup>, Mohammad Eftekhari<sup>1</sup>

<sup>1</sup> Research Center for Nuclear Medicine, Tehran University of Medical Sciences, Tehran, Iran

<sup>2</sup> Hasheminejad Kidney Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran

<sup>3</sup> Nuclear Medicine Department, Sina Hospital, Tehran University of Medical Sciences, Tehran, Iran

<sup>4</sup> Nuclear Medicine Department, Imam Hossein Hospital, Shahid Beheshti University of Medical Sciences, Tehran, Iran

(Received 12 September 2012, Revised 14 October 2012, Accepted 20 October 2012)

### ABSTRACT

**Introduction:** The aim of this study was to evaluate the effect of successful kidney transplantation (KT) on myocardial perfusion and left ventricular function by both qualitative (visual) interpretation and semiquantitative parameters, using myocardial perfusion scintigraphy with gated-single photon emission computed tomography (gated-SPECT) in patients suffering from end-stage renal disease.

**Methods:** From a total of 38 patients who were candidates of KT, twenty-six patients (16 female, 10 male, mean age: 47.5 yr, range: 24-64 yr) who had successful KT were included. Myocardial perfusion scintigraphy was performed by Gated Single Photon Emission Computed Tomography (Gated-SPECT) method, before and after surgery (mean: 24 months). Perfusion and function status was evaluated by qualitative and semiquantitative parameters.

**Results:** Our data showed qualitative evidence of perfusion and functional abnormality in pre-transplant scans as follows: Abnormal perfusion in left anterior descending (LAD), left circumflex (LCX) and right coronary artery (RCA) territories in 42.5%, 53.8% and 65.4% of cases, respectively; dilation in 57.7% and inhomogeneity of uptake in 53.8% of cases. However no statistically significant change was noted after transplantation, i.e. p value for all semiquantitative values including summed stress score (SSS), summed rest score (SRS) and summed difference score (SDS), summed motion score (SMS), summed thickening score (STS), ejection fraction (EF), end diastolic volume (EDV), end systolic volume (ESV), and stroke volume (SV) was greater than 0.05.

**Conclusion:** Renal transplantation may not have considerable long term effect on myocardial perfusion and function in patients with chronic renal failure. This could be due to either non-reversible myocardial changes or continuing effect of degrading factors on the myocardium.

**Key words:** Kidney transplantation; Myocardial perfusion; Myocardial function; Gated-SPECT

Iran J Nucl Med 2012;20(2):20-24

**Corresponding author:** Dr Davood Beiki, Research Center for Nuclear Medicine, Shariati Hospital, Tehran University of Medical Sciences, Tehran, Iran. E-mail: beikidav@tums.ac.ir

## INTRODUCTION

Chronic renal failure is a complex metabolic syndrome, and the uremic status can influence contractile function of myocardium [1]. It has also been shown that these patients might already have a large number of structural and functional peripheral vascular and cardiac abnormalities and the resultant myocardial ischemia may lead to left ventricular (LV) dysfunction that can persist even after the return of normal perfusion. This prolonged dysfunction is known as myocardial stunning [2].

It is suggested that kidney transplantation (KT) can resolve many of the cardiac abnormalities associated with chronic kidney failure [3,4].

Myocardial perfusion scintigraphy in gated SPECT mode is a sensitive and specific method that can effectively assess both cardiac perfusion and function [5,6]. We used this method before and after renal transplantation, to assess pre-transplantation myocardial perfusion and function in patients with end-stage renal disease, and subsequently to evaluate the probable effect of KT on improving perfusion and function status of myocardium.

## METHODS

Thirty-eight patients with chronic renal failure who were candidates for renal transplantation were enrolled in the study. Twelve patients were excluded because of unsuccessful transplantations or refusal to undergo the second scan after receiving transplant. A total of 26 patients (mean age: 47.5 yr, range: 24-64 yr), 16 female and 10 male, were included in the study.

For every patient myocardial perfusion scans, performed in gated SPECT mode before and after transplantation, were compared qualitatively and semi-quantitatively. The average interval between the two scans was 24 months and the average time between transplantation and the post-transplantation scan was 20 months.

This study was approved by the institutional ethics committee of Tehran University of Medical Sciences and written informed consent was obtained from all patients.

### *Image acquisition*

For myocardial perfusion imaging the patients were instructed to fast at least 4 hrs before the study. Possible interfering medications with dipyridamole study mainly xanthine containing drugs were stopped 48 hrs before the stress phase. Also caffeine containing foods and beverages were avoided for at least 24 hrs. A commercial sestamibi kit (AEOI,

Tehran, Iran) was used and the labeling and quality control procedures were performed according to the manufacturer's instructions. A dose of 666-814 MBq of Tc-99m sestamibi was given 4 min following the standard pharmacological stress with intravenous injection of 0.56 mg/kg dipyridamole over 4 min period. In the presence of dipyridamole side effects such as vertigo, chest pain, headache and electrocardiographic changes, 250 mg aminophylline was slowly injected intravenously 5 min after radiotracer injection. Single photon emission tomography (SPECT) with standard acquisition protocol was performed about 60 min after radiotracer injection, using a rotating, dual head gamma camera (Solus, ADAC, Milpitas, CA) equipped with a low-energy high resolution parallel hole collimator.

Patients were in a supine position during the image acquisition. Thirty-two azimuth images, 30 s/projection, were obtained in a 180-degree circular orbit, beginning from 45 degrees right anterior oblique to 135 degrees left posterior oblique with step and shoot acquisition on a 64×64 matrix and 38.5 cm detector mask (1.22 zoom). Rest images were obtained on the following day using the same imaging protocol. No attenuation correction was used in the imaging process.

### *Image analyses and interpretation*

Reconstruction of the images was carried out by Pegasys software (ADAC system). An expert nuclear physician used the cine-display of the rotating planar projections to assess sub-diaphragmatic activities, attenuations and patient motion to optimize the technical quality of the images.

The raw data were prefiltered by ramp and subsequently by Butterworth filters with frequency cut-off of 0.45 and order of 9.0. Also the data were semiquantitatively processed using Auto-QUANT software package (Cedars-Sinai Medical Center, Los Angeles, CA, USA) based on 20 segment analysis. Semiquantitative values, including EF, TID, LHR, SSS, SRS, SDS, SMS, STS, EDV, ESV and SV were also compared by statistical methods before and after renal transplantation.

The patients were stratified in four groups according to SSS, including normal ( $SSS \leq 3$ ), mildly abnormal ( $4 \leq SSS < 8$ ), moderately abnormal ( $9 \leq SSS < 11$ ), and severely abnormal ( $SSS \geq 12$ ).

Qualitative interpretation of scans, including homogeneity of uptake, visual perfusion status of myocardial walls, ventricular dilation and transient ventricular dilation (TID) was performed by two nuclear medicine physicians; in cases of disagreement a third opinion was also obtained.

**Statistical analysis**

We used SPSS statistical software (SPSS version 17.0) for statistical analysis. To compare semiquantitative values before and after renal transplantation, including EF, TID, LHR, SSS, SRS, SDS, SMS, STS, EDV, ESV and SV, wilcoxon signed rank test was used. To evaluate the qualitative data of pre and post transplantation measurements, Chi-Square test was used. A p-value of less than 0.05 was considered statistically significant.

**RESULTS**

Considering cardiac risk factors, from the total of 26 patients, 7 cases (26.9%) were hyperlipidemic, 14 (53.8%) hypertensive, 3 (11.5%) smoker and 5 (19.2%) were diabetic.

In assessing the perfusion status of pre-transplantation hearts by qualitative measure (visual interpretation), abnormal perfusion was noted with a frequency of 11 (42.5%) in LAD, 14 (53.8%) in LCX, and 17 (65.4%) in RCA territory. Visual dilation was noted in 15 (57.7%) and transient right ventricle visualization was seen in 6 (23.1%) of cases.

The post-transplantation scans were somehow more homogeneous than the pre-transplantation images,

though the value was considered insignificant ( $p=0.08$ ).

The difference in visual interpretation of perfusion in LAD territory (perfusing anterior and anteroseptal walls), LCX territory (perfusing anterolateral and inferolateral walls) and RCA territory (perfusing inferior and inferoseptal walls), before and after transplantation were insignificant (p values, 0.5, 0.1 and 1.0, respectively).

Regarding visual assessment, the frequency of left ventricular dilation, transient left ventricular dilation (TID), and transient right ventricle visualization were not different after transplantation compared to pre-transplantation studies ( $p=1.0$ ,  $p=0.1$ ,  $p=0.1$ , respectively). Semiquantitative data in pre- and post-transplantation scans are shown in Table 1.

In assessing SSS, from 21 patients who were within normal range ( $SSS \leq 3$ ), 19 patients (90.5%) remained in this group after transplantation, while only two patients were relocated as mildly abnormal group ( $8 \geq SSS > 4$ ). Mildly abnormal group ( $8 \geq SSS > 4$ ) included only one patient showing no change after transplantation. There were four patients in the severely abnormal group ( $SSS \geq 12$ ) from whom 3 patients remained in the same group, while one was changed to the medium risk group ( $11 \geq SSS > 9$ ).

**Table 1.** Comparison of semiquantitative perfusion and function data before and after renal transplantation.

Scintigraphic Parameter		Pre-transplantation	Post-transplantation	p value*
SSS	Median (range)	5 (0-22)	5 (0-21)	0.915
SDS	Median (range)	5 (0-7)	0 (0-11)	0.323
SRS	Median (range)	0 (0-18)	0 (0-21)	0.256
TID ratio	Mean (SD)	0.99±0.15	1.01±0.09	0.554
LHR	Mean (SD)	0.33±0.05	0.33±0.06	0.585
SMS	Median (range)	4.5 (0-69)	4 (0-70)	0.180
STS	Median (range)	0 (0-46)	0 (0-46)	0.311
ESV	Mean (SD)	33.50±23.05	33.72±23.36	0.482
EDV	Mean (SD)	77.04±34.26	78.28±36.35	0.530
EF	Mean (SD)	60.37±16.11	60.56±14.95	0.878

\* Wilcoxon Signed Ranks Test

SSS: summed stress score, SRS: summed rest score, SDS: summed difference score, TID ratio: transient ischemic dilation ratio, LHR: lung heart ratio, SMS : summed motion score, STS: summed thickening score, ESV: end systolic volume, EDV: end diastolic volume, EF: ejection fraction

## DISCUSSION

Chronic renal failure is a complex metabolic syndrome, and the uremic status can have adverse effect on myocardium [1]. This influence can be on myocardial blood supply or functional status of myocardium [2-4]. Uremia is likely to induce specific changes in the relaxation properties of the myocardium. These changes are responsible for the impaired diastolic function independent of blood pressure, degree of hypertrophy, and metabolic changes, suggesting the existence of a specific cardiomyopathy [7]. Prolonged exposure to uremic toxins has been shown to influence myocardial contractility. Although the mechanism of action of such toxins is not well established, several potentially negative inotropic and chronotropic factors have been verified in uremic plasma [8,9]. Exposure to these agents in long run can result in fibrosis of myocytes and eventually fatal cardiac event [10,11]. These patients might develop a specific cardiomyopathy, secondary to their specific metabolic, biochemical, hormonal, and even due to hemodynamic compromise (atrioventricular fistula) [12]. Cardiotoxic substances produced during uremia can adversely affect the myocardial function by slowing down its contraction and relaxation [13].

To evaluate cardiac perfusion and function in patients with chronic renal failure and to assess the effect of KT on cardiac status, electrocardiographically gated myocardial perfusion SPECT (GSPECT) method is the ideal method and was used in our study. This is a state-of-the-art technique for the combined evaluation of myocardial perfusion and left ventricular function within a single study. It is currently one of the most commonly performed cardiology procedures in nuclear medicine departments. Automation of the image processing and quantification has made this technique highly reproducible, practical and user friendly in the clinical setting. In patients with coronary artery disease, gating enhances the diagnostic and prognostic capability of myocardial perfusion imaging, providing incremental information over the perfusion data, and has shown potentials for myocardial viability assessment and sequential follow-up after therapy [5,14].

By this objective gateway method, myocardial perfusion, hypokinesia, dyskinesia, and functional parameters (left ventricular ejection fraction and systolic wall thickening) are well established and potentially useful to establish diagnostic, prognostic, and therapeutic indications [15].

Several studies have utilized Gated-SPECT as a valuable method in assessing prognosis after renal transplantation [4,16-19].

As renal transplantation causes significant improvement in azotemic condition, it is to be expected to induce improved cardiac status. Different studies have shown that renal transplantation can improve functional status of congestive heart failure [3,4,20,21], and increases survival [4]. Stokkel et al's study showed the systolic left ventricular dysfunction in terms of LVEF, wall motion, and wall thickening significantly improved 6 months after KT in patients with end-stage kidney disease [20]. Evaluation of heart function by echocardiography showed that after renal transplantation, myocardial hypertrophy was diminished and left ventricular systolic and diastolic function improved [3]. Parfrey showed improvement of LV systolic function, by an improvement in LV fraction shortening [21].

Our study however, didn't show such a trend. Although there was some evidence of improved perfusion after renal transplantation, the statistical evaluation did not show significant improvement in either perfusion or functional status of myocardium by qualitative or semiquantitative parameters. These parameters included visual interpretation of perfusion, dilation, transient RV visualization (qualitative assessment), TID, LHR, SSS, SRS, SDS, STS, SMS, EDV, ESV, SV, and EF (semiquantitative assessment), demonstrating p values persistently > 0.05. These findings could be due to long-standing uremic condition affecting cardiac perfusion and function causing irreversible changes of myocardium [2-4,7-11]. Another explanation for observation of coronary artery disease related abnormalities even after transplantation could be the possibility of increased risk of coronary artery calcification and coronary ischemia in renal transplant recipients [22]. This cardiovascular morbidity might also be caused by events related to surgical intervention and allograft function [23]. Also, the time interval after transplantation in our study (mean: 20 months) was longer than previous studies which showed functional improvement. We assumed the longer time interval after transplantation likely led to disappearance of the earlier improving cardiac effects explaining the findings in pre and post transplantation images in our study.

Also cardiac risk factors which might seem to have continuing influence on myocardium after transplantation were not interfering factors in our study considering their frequencies (hyperlipidemia: 26.9%, hypertension: 53.8%, smoking: 11.5% and diabetes: 19.2%).

The very similar findings in pre and post transplantation studies of our semiquantitative parameters (which are exclusively objective), while surprising, strongly confirmed our qualitative interpretations. More studies with larger sample size

are needed to further investigate these unexpected findings.

### CONCLUSION

Renal transplantation may not have considerable long term effect on myocardial perfusion and function in patients with chronic renal failure. This could be due to either non-reversible myocardial changes or continuing effect of degrading factors on the myocardium.

### Acknowledgements

This study has been funded and supported by Tehran University of Medical Sciences, Tehran, Iran; Grant no. 2759. Special thanks are also extended to the nuclear medicine technologists at our center for their technical help and data acquisition and also to Ghazal Haddad for her help in editing the manuscript.

### REFERENCES

- Hung J, Harris PJ, Uren RF, Tiller DJ, Kelly DT. Uremic cardiomyopathy--effect of hemodialysis on left ventricular function in end-stage renal failure. *N Engl J Med*. 1980 Mar 6;302(10):547-51.
- McIntyre CW. Haemodialysis-induced myocardial stunning in chronic kidney disease - a new aspect of cardiovascular disease. *Blood Purif*. 2010;29(2):105-10.
- Sahagún-Sánchez G, Espinola-Zavaleta N, Lafragua-Contreras M, Chávez PY, Gómez-Núñez N, Keirns C, Romero-Cardenas A, Pérez-Grovas H, Acosta JH, Vargas-Barrón J. The effect of kidney transplant on cardiac function: an echocardiographic perspective. *Echocardiography*. 2001 Aug;18(6):457-62.
- Wali RK, Wang GS, Gottlieb SS, Bellumkonda L, Hansalia R, Ramos E, Drachenberg C, Papadimitriou J, Brisco MA, Blahut S, Fink JC, Fisher ML, Bartlett ST, Weir MR. Effect of kidney transplantation on left ventricular systolic dysfunction and congestive heart failure in patients with end-stage renal disease. *J Am Coll Cardiol*. 2005 Apr 5;45(7):1051-60.
- Paul AK, Nabi HA. Gated myocardial perfusion SPECT: basic principles, technical aspects, and clinical applications. *J Nucl Med Technol*. 2004 Dec;32(4):179-87.
- Soman P. Gated SPECT myocardial perfusion scintigraphy: a multi-faceted tool for the evaluation of heart failure. *J Nucl Cardiol*. 2009 Mar-Apr;16(2):173-5.
- Facchin L, Vescovo G, Levedianos G, Zannini L, Nordio M, Lorenzi S, Caturelli G, Ambrosio GB. Left ventricular morphology and diastolic function in uraemia: echocardiographic evidence of a specific cardiomyopathy. *Br Heart J*. 1995 Aug;74(2):174-9.
- Vanholder R, Glorieux G, Lameire N. Uraemic toxins and cardiovascular disease. *Nephrol Dial Transplant*. 2003 Mar;18(3):463-6.
- Zoccali C, Bode-Böger S, Mallamaci F, Benedetto F, Tripepi G, Malatino L, Cataliotti A, Bellanuova I, Fermo I, Frölich J, Böger R. Plasma concentration of asymmetrical dimethylarginine and mortality in patients with end-stage renal disease: a prospective study. *Lancet*. 2001 Dec 22-29;358(9299):2113-7.
- Amann K, Breitbart M, Ritz E, Mall G. Myocyte/capillary mismatch in the heart of uremic patients. *J Am Soc Nephrol*. 1998 Jun;9(6):1018-22.
- Mall G, Huther W, Schneider J, Lundin P, Ritz E. Diffuse intermyocardial fibrosis in uraemic patients. *Nephrol Dial Transplant*. 1990;5(1):39-44.
- Drüeke T, Le Pailleur C, Meilhac B, Koutoudis C, Zingraff J, Di Matteo J, Crosnier J. Congestive cardiomyopathy in uraemic patients on long term haemodialysis. *Br Med J*. 1977 Feb 5;1(6057):350-3.
- McMahon AC, Vescovo G, Dalla Libera L, Wynne DG, Fluck RJ, Harding SE, Raine AE. Contractile dysfunction of isolated ventricular myocytes in experimental uraemia. *Exp Nephrol*. 1996 May-Jun;4(3):144-50.
- Go V, Bhatt MR, Hendel RC. The diagnostic and prognostic value of ECG-gated SPECT myocardial perfusion imaging. *J Nucl Med*. 2004 May;45(5):912-21.
- Bialostozky D, Leyva M, Villarreal T, Casanova JM, Pérez-Grovas H, Lemus P, Jiménez G, Vallejo E, Jiménez-Angeles L, Herrera J, Altamirano J. Myocardial perfusion and ventricular function assessed by SPECT and Gated-SPECT in end-stage renal disease patients before and after renal transplant. *Arch Med Res*. 2007 Feb;38(2):227-33.
- Patel AD, Abo-Auda WS, Davis JM, Zoghbi GJ, Deierhoi MH, Heo J, Iskandrian AE. Prognostic value of myocardial perfusion imaging in predicting outcome after renal transplantation. *Am J Cardiol*. 2003 Jul 15;92(2):146-51.
- Lewis MS, Wilson RA, Walker KW, Wilson DJ, Norman DJ, Barry JM, Bennett WM. Validation of an algorithm for predicting cardiac events in renal transplant candidates. *Am J Cardiol*. 2002 Apr 1;89(7):847-50.
- Hage FG, Smalheiser S, Zoghbi GJ, Perry GJ, Deierhoi M, Warnock D, Iskandrian AE, de Mattos AM, Aql RA. Predictors of survival in patients with end-stage renal disease evaluated for kidney transplantation. *Am J Cardiol*. 2007 Sep 15;100(6):1020-5.
- Garcheva M, Dzhorgova Iu, Nikolov G. Gated SPECT myocardial perfusion stress scintigraphy for evaluation of the ischemic risk in coronary artery disease. *Khirurgiia (Sofiia)*. 2007;(3):19-24.
- Stokkel M, Duchateau CS, Jukema W, de Fijter HW. Noninvasive assessment of left ventricular function prior to and 6 months after renal transplantation. *Transplant Proc*. 2007 Dec;39(10):3159-62.
- Parfrey PS, Harnett JD, Foley RN, Kent GM, Murray DC, Barre PE, Guttmann RD. Impact of renal transplantation on uremic cardiomyopathy. *Transplantation*. 1995 Nov 15;60(9):908-14.
- Seyahi N, Kahveci A, Cebi D, Altiparmak MR, Akman C, Uslu I, Ataman R, Tasci H, Serdengeci K. Coronary artery calcification and coronary ischaemia in renal transplant recipients. *Nephrol Dial Transplant*. 2011 Feb;26(2):720-6.
- Diaz G, O'Connor M. Cardiovascular and renal complications in patients receiving a solid-organ transplant. *Curr Opin Crit Care*. 2011 Aug;17(4):382-9.