

۲۰۱۱ Tl در اسکن پرفیوژن میوکارد با Reverse-Redistribution (مقاله مروری)

دکتر محمد افتخاری، دکتر رامین صادقی، دکتر ارمغان فرد اصفهانی، دکتر داود بیگی، دکتر بابک فلاحی، دکتر محسن ساغری

مؤسسه تحقیقات پزشکی هسته‌ای، بیمارستان دکتر شریعتی، دانشگاه علوم پزشکی و خدمات بهداشتی-درمانی تهران

چکیده

نمای reverse redistribution در اسکن پرفیوژن میوکارد به معنی کاهش اکتیویته یک ناحیه از میوکارد در مرحله استراحت نسبت به مرحله استرس است. در مورد علت و اهمیت بالینی این پدیده بحث زیادی شده است، ولی هنوز هم نظرات متفاوتی در این مورد اظهار می‌شود. در مورد مکانیزم این پدیده نظر غالب بر این است که خروج زودرس رادیودارو علت آن می‌باشد. در مورد شرایط میوکارد به خصوص زنده بودن (viability) هم تقریباً اتفاق نظر وجود دارد و اکثر مطالعات به زنده بودن نواحی مبتلا به reverse redistribution اشاره دارند، در حالیکه مطالعات مختلف در مورد پیش‌آگهی نواحی مبتلا به RR نظرات متفاوتی ابراز کرده‌اند.

در این مقاله سعی شده است مطالعات مختلف در مورد پدیده reverse redistribution را با رادیوداروی ²⁰¹Tl بررسی شده و در نهایت یک نتیجه گیری کلی که از نظر بالینی بتواند جهت متخصصان پزشکی هسته‌ای راهگشایی باشد ارائه گردد.

واژه‌های کلیدی: reverse redistribution، اسکن پرفیوژن میوکارد، ²⁰¹Tl

بسته به نرمال یا غیر نرمال بودن ناحیه RR در تصاویر استرس، نمای RR به دو قسمت تقسیم می‌گردد RR-A و RR-B (RR-4-6). در نوع اول ناحیه RR در تصاویر استرس نرمال است، ولی در نوع دوم کاهش برداشت رادیودارو در مرحله استرس هم دیده می‌شود.

نواحی A-RR نسبت به B-RR موج Q کمتر نشان میدهند و به Revascularization بeter جواب می‌دهند. Pace و همکاران پیشنهاد کرده اند که نواحی دیس سینرژیک با نمای RR-A باید زنده تلقی شوند، در حالیکه در مورد زنده بودن نواحی RR-B باید با احتیاط برخورد کرد (۴).

تاریخچه

اولین بار RR توسط Tanasecu و همکاران در سال ۱۹۷۹ گزارش شد (۷). پس از آن Hecht و همکاران در سال ۱۹۸۱ لفظ Reverse-Redistribution را در مورد این پدیده به کار برداشتند (۸). پس از معمول شدن روش

مقدمه

یکی از مشکلاتی که هر متخصص پزشکی هسته‌ای در گزارش اسکن پرفیوژن میوکارد با آن رویرو است نمای (RR) Reverse-Redistribution است. با وجود گذشت سالها از شروع استفاده از تالیم در اسکن پرفیوژن میوکارد، هنوز هم عقاید بسیار متفاوتی در مورد RR موجود است. در این مقاله سعی شده است تا با بررسی مقالات مختلف به یک جمع بندی نهایی در این مورد دست یابیم.

تعريف

کاهش اکتیویته در یک ناحیه از میوکارد به میزان حداقل ۱۰٪ در تصاویر استراحت نسبت به استرس مشخصه RR است (۱ و ۲).

به نظر می‌رسد اگر RR در تصاویر پس از تزریق مجدد ۳ ساعته دیده شود، نشان دهنده ضایعه شدیدتری در میوکارد نسبت به وجود آن در تصاویر ۲۴ ساعته است (۳).

خصوص پس از MI و بدنبال درمان ترومبولیتیک دیده می شود. نواحی RR در واقع دچار خایعه ناکامل میوکارد شده اند و قادر به نگهداری^{۲۰۱} TI به اندازه مناطق نرمال نیستند. به خصوص اگر رگ ترومبوزه کاملاً باز شده باشد، به علت افزایش شدید جریان خون به نواحی RR این حالت بارزتر است(۲۷).

(۲) برداشت تأخیری و خروج تسریع شده: این مکانیزم به خصوص توجیه کننده RR در تصاویر تزریق مجدد است(۲۶).

(۳) آرتکت ها: عدم دقّت در مراحل تصویربرداری و process اسکن پرفیوژن میوکارد می تواند عامل ایجاد RR باشد. به عنوان مثال RR در نواحی آپیکال و قدامی قلب می تواند به علت تغییر محل پستان در بین دو مرحله استرس و استراحت باشد. حتی گزارش شده است که پراکنش اکتیویته از نواحی اطراف به خصوص احشاء شکمی می تواند اکتیویته ناحیه تحتانی را در استراحت افزایش دهد و موجب RR در نواحی قدامی و آپیکال گردد(۲۸). Brown و همکاران و Lear و همکاران به طور جداگانه تفریق زمینه را به عنوان علت RR ذکر کرده اند، که البته فقط در مورد تصاویر planar صدق می کند(۲۹و۳۰). به هر حال در بررسی RR حتماً باید آرتکت ها را در نظر داشت و ابتدا آنها را رد کرد(۲۸).

(۴) واریاسیون طبیعی در خروج^{۲۰۱} TI در افاد با کرونر طبیعی: Kaul و همکاران این پدیده را گزارش کرده اند(۱۰).

اهمیت بالینی RR در اسکن پرفیوژن میوکارد:

آیا RR نشان دهنده تنگی عروق کرونر خونرساننده است؟

اطلاعات بسیار متناقضی در این مورد ارائه شده است. Tanasescu و همکاران نشان داده اند که نواحی RR توسط عروق کرونر نرمال یا با تنگی خفیف Weiss و همکاران نیز نشان داده اند که نمای RR پس از MI نشانه باز بودن عروق کرونر خونرساننده آن نواحی است(۳۱و۱۲).

از طرف دیگر Hecht و همکاران، RR را نشانه درگیری شدید عروق کرونر همان نواحی می دانند(۸).

تزریق مجدد در اسکن پرفیوژن میوکارد اولین بار Dilsizian و همکاران در سال ۱۹۹۲ این پدیده را شرح دادند(۹).

در چه مواردی دیده میشود؟

- (۱) افراد با عروق کرونر نرمال(۱۰).
- (۲) نواحی مبتلا به CAD مزمن: فراوانی RR در این نواحی کم است. Hecht و همکاران و Soufer و همکاران این میزان را به ترتیب ۷٪/۱۵٪ ذکر کرده اند(۱۱و۸).

(۳) نواحی مبتلا به MI پس از Re-perfusion فراوانی RR در این نواحی بسیار بالا گزارش شده است. حتی Weiss و همکاران این میزان را تا حد ۷۵٪ ذکر کرده است(۱۲).

(۴) سایر موارد: در مقالات مختلف و گزارش موردهای متعددی، RR گزارش شده است مانند: هیپرتروفی بطنی(۱۳)، کاردیومیوپاتی ها(Sarcoidosis)(۱۴)، شاگاس(۱۵) و ایسکمیک(۱۶)، آریتمی ها(WPW)(۱۷)، پل میوکاردی(۱۸)، آنژیوپلاستیک(۱۹)، کاوساکی(۲۰)، پیوند قلب(۲۱و۲۲)، سندروم X(۲۳)، MI غیر ترانس سورال(۲۴) و بیماری های کلژن- و اسکولار(۲۵).

مکانیزم ایجاد RR

در مورد مکانیزم ایجاد RR بحث فراوانی شده است و مطالعات مختلف مکانیزم های متفاوتی را در این مورد مطرح کرده اند.

Beygui و همکاران ۳ مرحله را در جذب سلوی^{۲۰۱} TI ذکر می کنند: مرحله اولیه واسکولا، مرحله بینایینی (interstitial) و مرحله سلوی نهایی. اختلال در هر کدام از این مراحل موجب ایجاد نمای غیر طبیعی در اسکن می گردد. به طور کلی ورود^{۲۰۱} TI به سلوی نیازمند جریان خون نرمال و متابولیزم سلوی مناسب است و خروج آن بستگی به متابولیزم سلوی و تمامیت (integrity) غشاء سلوی دارد(۲۸و۲۶). بر این اساس مکانیزم های RR به صورت زیر پیشنهاد شده اند:

- (۱) افزایش خروج^{۲۰۱} TI ثانویه به افزایش جریان خون به ناحیه نجات یافته از انفارکتوس میوکارد: این حالت به عنوان علت اصلی RR مطرح است و به

اختلال حرکت دیواره قلب در نواحی RR پس از CABG بهبود می‌یابد(۲۷).

تمام مطالعات فوق در مورد بیمارانی بوده است که یک پروسه درمانی اعم از درمان ترومبولیتیک یا CABG بر روی آنها انجام شده است. سؤال بالینی مهم این است که آیا نمای RR در بیماران مبتلا به ایسکمی مزمن میوکارد می‌تواند نشانه‌زنده بودن باشد.

در یک مطالعه مقایسه‌ای با $^{18}\text{FDG-PET}$ ^{۱۸} در بیماران مبتلا به ایسکمی مزمن میوکارد Marin-neto و همکاران نتیجه گرفته اند که حدود ۵۰٪ ناحیه از ۳۶ ناحیه) از نواحی RR دارای زنده بودن هستند(۱۵). در یک مطالعه بسیار دقیق توسط Soufer و همکاران نواحی RR در بیماران دچار CAD مزمن با $^{13}\text{NH}_3$ ^{۱۳} مقایسه شده اند. کلاً ۷۷٪(۳۶ ناحیه از ۵۰ ناحیه) از این نواحی زنده بوده اند. ۵۰٪ نواحی دچار اختلال حرکت دیواره و ۹۰٪ نواحی بدون اختلال حرکت زنده بودند و بنابراین وجود اختلال حرکتی در نواحی RR نشانه عدم وجود زنده بودن نیست. در همین مطالعه بیماران پی گیری ۱۴ ماهه شدند و ۵۰٪ افراد دارای RR دچار حادثه قلبی شدند(در همان نواحی دارای RR). در نتیجه وجود RR در اسکن پر فیوژن میوکارد باید جدی تلقی شده و هر چه سریعتر درمان شود(۱۱).

در مطالعه دیگر توسط Ohte و همکاران نواحی RR با $^{18}\text{FDG-PET}$ ^{۱۸} مقایسه شده اند و مؤلفان نتیجه گرفته اند که مصرف گلوکز در نواحی RR کاهاش می‌یابد ولی نه به اندازه نواحی مبتلا به اسکار(۴۱).

RR بیانگر پیش آگهی خوب است یا بد؟

در این مورد هم اطلاعات ضد و نقیض فراوان است. با این وجود اکثر مطالعات نشان دهنده پیش آگهی خوب هستند(۴۲-۴۶).

De Sutter و همکاران بیان می‌دارند که وجود RR در اسکن پر فیوژن میوکارد بیماران دچار MI پس از آنژیوپلاستی نشانه پیش آگهی خوب در یکسال آینده است.^{۴۷} مطالعات متعددی نیز همین نظر را تأیید می‌کنند(۴۷-۴۸ و ۳۵-۳۱).

Sakata و همکاران در یک مطالعه ۵۰ بیمار را پس از انجام ترومبولیز تحت اسکن پر فیوژن میوکارد قرار

نستایج مطالعات Koliakos و همکاران Pace و همکاران نیز با این مسئله هماهنگی داشته و RR در نواحی دارای عروق کرونر بیمار دیده شده است(۳۲ و ۳۶). گروه دیگری از مطالعات هیچ ارتباطی بین RR و شدت درگیری عروق کرونر نیافته اند. به عنوان مثال Silberstein و همکاران Lee.J.K و همکاران نشان داده اند که RR می‌تواند هم در عروق کرونر بیمار و هم در عروق کرونر نرمال دیده شود(۳۳ و ۳۴).

آیا RR به معنی برقراری جریان خون ناحیه‌ای میوکارد پس از درمان ترومبولیتیک است؟

در این مورد نظرات تقریباً یکسان هستند. کلاً عقیده بر این است که نواحی RR پس از درمان ترومبولیتیک یا عروق کرونر باز دارند یا توسط عروق کلاترال به خوبی مشروب می‌شوند(۳۱ و ۳۶).

آیا RR به معنی وجود viability است؟

در این مورد هم اتفاق نظر وجود دارد و عقیده بر این است که در نواحی RR مخلوطی از نسج زنده و اسکار وجود دارد(۳۵ و ۱۲).

مطالعات فراوانی در مورد وضعیت نواحی RR پس از درمان ترومبولیتیک انجام شده است و تقریباً تمام آنها به نفع وجود زنده بودن در نواحی RR هستند. به عنوان مثال Touchstone و همکاران نشان داده اند که نواحی RR پس از درمان ترومبولیتیک بهبود قابل توجه در عملکرد سیستولی می‌یابند(۳۶). Yamagishi و همکاران در دو مطالعه جداگانه نشان داده اند که نواحی RR ۱ هفته الی ۲ ماه پس از MI به تدریج بهبود عملکرد پیدا می‌کنند(۳۷-۳۹).

در مطالعه دیگر Langer و همکاران نشان دادند که نواحی RR پس از مصرف نیتروگلیسیرین زیرزبانی نسبت به نواحی ثابت یا دارای نقص برگشت پذیر بهبود عملکرد بیشتری پیدا می‌کنند(۳۱).

Faraggi و همکاران هم RR را نشانه زنده بود نمی‌دانند و ۸ روز پس از درمان ترومبولیتیک را بهترین زمان برای پیش بینی بهبود عملکرد میوکارد در نظر گرفته است(۴۰).

در تنها مطالعه انجام شده در مورد اهمیت RR پس از CABG و همکاران Watarida نشان داده اند که

نتیجه‌گیری نهائی

Berman و Maddahi اهمیت RR را دراسکن پرفیوژن میوکارد با ^{201}TI به صورت زیر خلاصه کرده‌اند (۵۲):

- (۱) افراد با احتمال کم CAD: RR در این افراد به احتمال زیاد یک واریاسیون نرمال است یا به علت آرنفکت‌ها است، که مورد آخر حتماً باید در تمام موارد RR ابتدا در نظر گرفته شده و رد شود.
- (۲) در افراد مبتلا به CAD مزمن یا بدنبال درمان تروموبلیتیک اخیر: RR در این افراد نشانه وجود مخلوط‌طی از نسج زنده و اسکار است که توسط عروق باز یا کلاترال‌ها به خوبی خونرسانی می‌شوند. پیش‌آگهی در این افراد معمولاً خوب است.
- (۳) در افرادی که RR شدید نشان می‌دهند استفاده از $^{18}\text{FDG-PET}$ برای تعیین احتمال حوادث عروقی آینده می‌تواند بسیار مفید باشد.

داده‌اند و نتیجه می‌گیرند که نواحی RR یا ثابت نسبت به نواحی برگشت پذیر میزان کمتری از MI مجدد دارند (۴۹).

مطالعاتی نیز وجود دارند که نشانه پیش‌آگهی بد در افرادی است که نمای RR را در اسکن پرفیوژن میوکارد نشان می‌دهند. Wackers و همکاران نشان داده‌اند که وجود RR پس از تروموبلیز باعث افزایش بروز عوارض قلبی-عروقی می‌شود (۵۰). همان طور که قبل از ذکر شد وجود RR شدید که در بررسی با $^{18}\text{FDG-PET}$ شواهدزنده بودن را نشان دهد، همراه با افزایش حوادث قلبی است (۱۱). Faragasso و همکاران در یک گزارش مورد جالب دو بیمار را معرفی کرده‌اند که به علت درد قلبی اسکن پرفیوژن میوکارد با ^{201}TI شده‌اند و نمای RR داشته‌اند و هر دوی آنها پس از مدتی در همان ناحیه RR دچار شده‌اند (۵۱).

منابع

- 1) Sciagra R, Pupi A, Pellegrini M, Matteini M, Santoro GM, Bisi G and Fazzini PF; Prediction of post-revascularization functional recovery of asynergic myocardium using quantitative 201TI rest-redistribution tomography: Has the reverse redistribution pattern an independent significance? Eur J Nucl Med 1998; 25(6): 594-600
- 2) Pace L, Perrone FP, Mainenti PP, Prastaro M, Cuocolo A, Varrone A, Vezzuto P, Crisci T, Soricelli A, Piscione F, Chiariello M and Salvatore M; Effects of myocardial revascularization on regional 201TI uptake and systolic function in regions with reverse redistribution on tomographic 201TI imaging at rest in patients with chronic coronary artery disease. J Nucl Cardiol 1998; 5(2): 153-160
- 3) Ohte N, Hashimoto T, Iida A, Narita N and Akita S; Extent of myocardial damage in regions with reverse redistribution at 3 h and at 24 h on 201TI SPET: evaluation based on regional myocardial oxidative metabolism. Nucl Med Commun 1998; 19(11):1081-7
- 4) Pace L, Cuocolo A, Marzullo P, Nicolai E, Gimelli A, De Luca N, Ricciardelli B and Salvatore M; Reverse redistribution in resting 201TI myocardial scintigraphy in chronic coronary artery disease: an index of myocardial viability. J Nucl Med 1995; 36(11):1968-73
- 5) Pace L, Cuocolo A, Nicolai E, Imbriaco M, Maurea S, Nappi A, Ricciardelli B and Salvatore M; Reverse redistribution in 201TI stress-redistribution myocardial scintigraphy, Effect of rest reinjection. Clin Nucl Med 1994; 19(11):956-61
- 6) Pace L, Cuocolo A, Maurea S, Nicolai E, Imbriaco M, Nappi A, Morisco C, Chiariello M, Trimarco B and Salvatore M; Reverse

- redistribution in resting 201Tl myocardial scintigraphy in patients with coronary artery disease: relation to coronary anatomy and ventricular function. *J Nucl Med* 1993; 34(10):1688-92
- 7) Tanasescu D, Berman D and Staniloff H; Apparent worsening of 201Tl myocardial defects during redistribution - what does it mean? *J Nucl Med* 1979; 20:688
- 8) Hecht HS, Hopkins JM, Rose JG, Blumfield DE and Wong M; Reverse redistribution: worsening of 201Tl myocardial images from exercise to redistribution. *Radiology* 1981; 140:177-181
- 9) Dilsizian V and Bonow RO; Differential uptake and apparent 201Tl washout after thallium reinjection: options regarding early redistribution imaging before reinjection and late redistribution imaging after reinjection. *Circulation* 1992; 85:1032-1038
- 10) Kaul S, Chesler DA, Newell JB, Pohost GM, Okada RD and Boucher CA; Regional variability in the myocardial clearance of 201Tl and its importance in determining the presence or absence of coronary artery disease. *J Am Coll Cardiol* 1986; 8:95-100
- 11) Soufer R, Dey HM, Lawson AJ, Wackers FJT and Zaret BL. Relationship between reverse redistribution on planar thallium scintigraphy and regional myocardial viability: a correlative positron emission tomography study. *J Nucl Med* 1995; 36:180-187
- 12) Weiss AT, Maddahi J, Lew AS, Shah PK, Ganz W, Swan JH and Berman DS; Reverse redistribution of 201Tl: a sign of nontransmural myocardial infarction with patency of the infarct-related coronary artery. *J Am Coll Cardiol* 1986; 7:61-67
- 13) Candell Riera J and Ortega Alcalde D; Reverse redistribution pattern of 201Tl stress test in subjects with normal coronary angiograms. *J Nucl Med* 1986; 27:1377
- 14) Fields CL, Ossorio MA, Roy TM, Denny DM and Varga DW; 201Tl scintigraphy in the diagnosis and management of myocardial sarcoidosis. *South Med J* 1990; 83:339-342
- 15) Marin-Neto JA, Marzullo P and Marcassa C; Myocardial abnormalities in chronic Chagas' disease as detected by 201Tl scintigraphy. *Am J Cardiol* 1992; 69:780-784
- 16) Noah MS; Non-invasive evaluation of the myopathy of coronary artery disease. *Brit J Clin Pract* 1989 Jul; 43(7):249-51
- 17) Nil T, Nakashima Y, Nomoto J, Hiroki T, Oshima F and Arakawa K; Normalization of reverse redistribution of 201Tl with procainamide pretreatment in Wolff-Parkinson-White syndrome. *Clin Cardiol* 1991; 14:269-72
- 18) Ho YL, Wu CC, Yen RF, Hung SR, Chen MF and Huang PJ; Comparison of ischemic patterns in myocardial bridge and syndrome X: Evaluation by dobutamine stress echocardiography and stress 201Tl SPECT. *J Formos Med Assoc* 2001; 100(2): 83-88
- 19) Nakano A, Lee JD, Shimizu H, Tsubokawa A, Uzui H, Tsuchida T, Takahashi N, Yonekura Y and Ueda T; Clinical significance of reverse redistribution on resting 201Tl imaging in patients with vasospastic angina. *Ann Nucl Med* 2001; 15(1): 65-68
- 20) Tsai CH, Lee JK, Kao CH, Chang JS and Shih WJ; Kawasaki disease evaluated by two-dimensional echocardiogram and dipyridamole (201)Tl-chloride myocardial SPET. *Nucl Med Commun* 1997; 18(5): 412-418
- 21) Puskas C, Kosch M, Kerber S, Jonas M,

- Weyand M, Breithardt G, Scheld HH and Schober O; Progressive heterogeneity of myocardial perfusion in heart transplant recipients detected by ^{201}Tl myocardial SPECT. *J Nucl Med* 1997; 38(5): 760-765
- 22)Kerber S, Puschkas C, Jonas M, Janssen F, Heinemann VO, Kosch M, Deng MC, Schober O, Scheld HH and Breithardt G; Can ^{201}Tl myocardial SPECT abnormalities in orthotopic heart recipients be explained by coronary vessel wall alterations assessed by intravascular ultrasound? *Int J Cardiol* 1996; 57(1): 91-96
- 23)Fragasso G, Rossetti E, Dosio F, Gianolli L, Pizzetti G, Cattaneo N, Fazio F and Chierchia SL; High prevalence of the ^{201}Tl reverse redistribution phenomenon in patients with syndrome X. *Eur Heart J* 1996; 17(10): 1482-1487
- 24) Choe W, Kwan J and Kim S; Reverse redistribution: revisited with myocardial contrast echocardiography. *Ann Nucl Med* 2000 Feb; 14(1):63-70
- 25)Ishida R, Murata Y, Sawada Y, Nishioka K and Shibuya H; ^{201}Tl myocardial SPET in patients with collagen disease. *Nucl Med Commun* 2000; 21(8): 729-734
- 26)Beygui F, Feuvre C, Maunoury C, Helft G and Metzger JF; Coronary Vasodilator Reserve: A clue to the explanation of ^{201}Tl redistribution patterns early after successful primary stenting for acute myocardial infarction. *J Am Coll Cardiol* 2002; 40(5): 300-305
- 27)Watarida S, Onoe M, Sugita T, Tabata R, Katsuyama K, Nakajima Y, Nojima T, Yamamoto R, Matsuno S and Mori A; Clinical significance of Reverse Redistribution phenomenon after coronary bypass grafting. *Ann Thorac Surg* 1995; 59:1528-33
- 28)Faraggi. M; The continuing story of ^{201}Tl reverse redistribution: Reverse Redistribution is still alive, but is the myocardium still viable? *J Nucl Med* 2002;43(5):628-631
- 29) Brown KA, Benoit L, Clements JP and Wackers FJ; Fast washout of ^{201}Tl from area of myocardial infarction: possible artifact of background subtraction, *J Nucl Med* 1981; 28:945-949
- 30)Lear JL, Raff U and Jain R. Reverse and pseudo redistribution of ^{201}Tl in healed myocardial infarction and normal and negative ^{201}Tl washout in ischemia due to background oversubtraction. *Am J Cardiol* 1988; 62:543-550
- 31)Langer A, Burns RJ, Freeman MR, Liu P, Morgan CD, Wilson R and Armstrong PW; Reverse redistribution on exercise thallium scintigraphy: relationship to coronary patency and ventricular function after myocardial infarction. *Can J Cardiol* 1992; 8:709-715
- 32)Koliakos G, Doumas A, Altas D and Louridas G; The clinical significance of reverse redistribution in Tl^{201} cardiac SPET. *Int J Cardiovasc Imag* 2001 Feb; 17(1):29-35
- 33)Silberstein EB and DeVries DF; Reverse redistribution phenomenon in ^{201}Tl stress tests: angiographic correlation and clinical significance. *J Nucl Med* 1985; 26:707-710.
- 34)Lee JK and Kao CH; Reverse redistribution during dipyridamole ^{201}Tl myocardial single photon emission computed tomography in coronary artery disease. *Kaohsiung J Med Sci* 1996 Nov; 12(11):630-633
- 35)Fukuzawa S, Ozawa S, Nobuyoshi M and Inagaki Y; Reverse redistribution on ^{201}Tl SPECT images after reperfusion therapy for acute myocardial infarction: possible mechanism and prognostic implications.

- Heart Vessels 1992; 7:141-147 .
- 36)Touchstone DA, Beller GA, Nygaard TW, Watson DD, Tedesco C and kaul S; Functional significance of predischarge exercise 201Tl findings following intravenous streptokinase therapy during acute myocardial infarction. Am Heart J 1988; 116:1500-1507.
- 37)Yamagishi H, Itagane H and Akioka K; Implications of delayed image on simultaneous 201Tl single-photon emission computed tomography in patients with acute myocardial infarction. Jpn Circ J 1993; 57:27-36.
- 38)Yamagishi H, Itagane H, Akioka K, Ohmura T, Iida H, Tahara A, Toda I, Teragaki M, Takeuchi K and Takeda T; Clinical significance of reverse redistribution on 201Tl single-photon emission computed tomography in patients with acute myocardial infarction. Jpn Circ J 1992; 56(11): 1095-1105
- 39)Umamoto I, Sugihara H, Harada Y, Sawada T, Matsumuro A, Matsubara K, Shiga K, Nakagawa T, Oonishi K, Nakamura T, Azuma A, Kohno Y, Katsume H, Nakagawa M and Miyao K; The clinical significance of reverse redistribution of 201Tl SPECT at rest in the 1st month after the onset of acute MI. Kakuigaku 1991; 28(7): 749-757
- 40)Faraggi M, Karila CD, Brochet E, Lebtahi R, Czitrom D, Feldman LJ, Assayag P, Doumit D, Steg G and Le Guludec D; Relationship between resting 201Tl reverse redistribution, microvascular perfusion, and functional recovery in acute myocardial infarction. J Nucl Med 2000 Mar; 41(3):393-399
- 41)Ohte N, Hashimoto T, Banno T, Narita H, Kobayashi K, Akita S and Fujinami T; Clinical significance of reverse redistribution on 24-hour delayed imaging of exercise 201Tl myocardial SPECT: comparison with myocardial fluorine-18-FDG-PET imaging and left ventricular wall motion. J Nucl Med 1995 Jan; 36(1):86-92
- 42)De Sutter J, Van de Wiele C, Dierckx R, Gheeraert P, De Buyzere M and Taeymans Y; Reverse redistribution on 201Tl single-photon emission tomography after primary angioplasty: a one-year follow-up study. Eur J Nucl Med 1999 Jun; 26(6):633-639
- 43)Sciagra R, Pupi A, Pellegrini M, Matteini M, Santoro GM, Bisi G and Fazzini PF; Prediction of post-revascularization functional recovery of asynergic myocardium using quantitative 201Tl rest-redistribution tomography: has the reverse redistribution pattern an independent significance? Eur J Nucl Med 1998 Jun; 25(6):594-600
- 44)Pace L, Perrone-Filardi P, Mainenti PP, Prastaro M, Cuocolo A, Varrone A, Vezzuto P, Crisci T, Soricelli A, Piscione F, Chiariello M and Salvatore M; Effects of myocardial revascularization on regional 201Tl uptake and systolic function in regions with reverse redistribution on tomographic 201Tl imaging at rest in patients with chronic coronary artery disease. J Nucl Cardiol 1998 Mar-Apr; 5(2):153-160
- 45)Roelants VA, Vanoverschelde JL, Vander Borght TM and Melin JA; Reverse redistribution on exercise-redistribution (201)Tl SPECT in chronic ischemic dysfunction: predictive of functional outcome after revascularization? J Nucl Med 2002 May ; 43(5):621-627
- 46)Hirata K, Yamagishi H, Tani T, Sakanoue Y, Akioka K, Takeuchi K, Yoshikawa J and Ochi H; Reverse redistribution of 201Tl myocardial single photon emission

- tomography and contractile reserve. *Jpn Circ J* 2000 May; 64(5):345-351
- 47) Sridhara BS, Dudzic E, Basu S, Senior R and Lahiri A; Reverse redistribution of 201Tl represents a low-risk finding in thrombolysed patients following myocardial infarction. *Eur J Nucl Med* 1994 Oct; 21(10):1094-7
- 48) Dudzic EM, Sridhara BS and Lahiri A; Reverse redistribution: fact or fiction? *Eur J Nucl Med* 1994 May; 21(5):449-53
- 49) Sakata K, Yoshida H and Ono N; Implications of delayed image on simultaneous 201Tl/technetium-99m pyrophosphate dual emission computed tomography early after acute myocardial infarction. *Jpn Circ J* 1993; 57:27-36.
- 50) Wackers FJT, Zaret BL, Chatman B, Wassermann A and Thompson B; The prognostic significance of reverse redistribution in 201Tl stress testing after thrombolytic therapy for acute infarction. *J Am Coll Cardiol* 1992; 22:100-105
- 51) Fragasso G, Chierchia SL, Pizzetti G, Dosio F and Fazio F; Reverse redistribution of 201Tl heralding the development of myocardial infarction: description of two cases. *J Nucl Biol Med* 1994 Sep; 38(3):515-7
- 52) Maddahi J and Berman DS; Reverse redistribution of 201Tl. *J Nucl Med* 1995; 36:1019-1021.